

This is a repository copy of *Transcription factors in eukaryotic cells can functionally regulate gene expression by acting in oligomeric assemblies formed from an intrinsically disordered protein phase transition enabled by molecular crowding*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/130472/>

Version: Accepted Version

---

## Article:

Leake, Mark Christian orcid.org/0000-0002-1715-1249 (2018) Transcription factors in eukaryotic cells can functionally regulate gene expression by acting in oligomeric assemblies formed from an intrinsically disordered protein phase transition enabled by molecular crowding. *Transcription*. pp. 298-306.

<https://doi.org/10.1080/21541264.2018.1475806>

---

## Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

## Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Transcription

Transcription factors in eukaryotic cells can functionally regulate gene expression by acting in oligomeric assemblies formed from an intrinsically disordered protein phase transition enabled by molecular crowding

--Manuscript Draft--

Manuscript Number:	KTRN-2018-0012R1
Article Type:	Point-of-View
Keywords:	gene expression; transcription factors; single-molecule; super-resolution; cell signaling; intrinsically disordered protein; phase transition; molecular crowding; fluorescent protein
Corresponding Author:	Mark Leake  UNITED KINGDOM
First Author:	Mark Leake
Order of Authors:	Mark Leake
Manuscript Region of Origin:	UNITED KINGDOM
Abstract:	High-speed single-molecule fluorescence microscopy in vivo shows that transcription factors in eukaryotes can act in oligomeric clusters mediated by molecular crowding and intrinsically disordered protein. This finding impacts on the longstanding puzzle of how transcription factors find their gene targets so efficiently in the complex, heterogeneous environment of the cell.

<sup>a</sup> Biological Physical Sciences Institute, Departments of Physics and Biology, University of York, York YO10 5DD, UK

Corresponding author: Prof Mark Leake, Biological Physical Sciences Institute, Departments  
of Physics and Biology, University of York, York YO10 5DD, UK. Tel: +44 (0)1904322697.  
Email: mark.leake@york.ac.uk. Orcid ID: <http://orcid.org/0000-0002-1715-1249>.

**Keywords:** gene expression; transcription factors; single-molecule; super-resolution; cell signaling; intrinsically disordered protein; phase transition; molecular crowding; fluorescent protein

## Abstract

High-speed single-molecule fluorescence microscopy in vivo shows that transcription factors in eukaryotes can act in oligomeric clusters mediated by molecular crowding and intrinsically disordered protein. This finding impacts on the longstanding puzzle of how transcription factors find their gene targets so efficiently in the complex, heterogeneous environment of the cell.

## Introduction

Cells regulate gene expression through binding of transcription factors (TFs) to promoters to turn gene expression on or off (1, 2). Simulations show that the time it takes for TFs to find their targets through pure 3D diffusion alone is ~six orders of magnitude larger than what is observed experimentally (3). Hypotheses to explain this observation have included TF heterogeneous mobility comprising a combination of free 3D diffusion combined with sliding and hopping on the DNA plus longer jumps between different DNA strands called intersegment transfer (4–6). In eukaryotic cells, TF localization fluctuates, often between cytoplasm and nucleus (7). Although it has been observed that promoters can pool on the genome in clusters (8) it has not previously been seen that TFs themselves act in clusters, but instead are largely assumed to act as single molecules. Simulations which embody diffusion and binding suggest that multivalent TFs could, in principle, facilitate intersegment transfer (9). Previously, single-molecule fluorescence microscopy has been used to study TF localization in living cells across a range of model organisms, including bacteria, yeast and multi-cellular organisms (10–16). Many studies suggest complexities in diffusion and binding (4, 12, 15, 17, 18) which may include intersegmental transfer (4, 17, 18). However, until now, the direct experimental evidence for intersegmental transfer has been limited.

Many of the important features of gene expression control in eukaryotes are exemplified in the model unicellular microorganism *Saccharomyces cerevisiae* (budding yeast). In particular, its glucose sensing pathway presents an experimentally tractable system to study gene regulation. Here, control of gene expression is achieved by TFs which include the Zn finger DNA binding protein Mig1 (19) that acts to repress expression from targets including *GAL* genes involved in glucose metabolism (20). Mig1 localizes towards the nucleus if the extracellular glucose concentration is increased (21), correlated to its own dephosphorylation by a protein called Snf1 (22, 23).

In recent investigations from my own group (24) the spatiotemporal dynamics and kinetics of gene regulation in live *S. cerevisiae* cells, using its glucose sensing pathway as a model for signal transduction, was explored using physics methods which enable the understanding of the processes of life one molecule at a time (25, 26), employing ‘single-molecule optical proteomics’ tools (27). The combination of these advanced light microscopy with genetics techniques has previously enabled valuable insights into the activities of several other processes for low copy number proteins (28) in both unicellular organisms and single cells from more complex multicellular organisms (29). These single-molecule/cell and super-resolution microscopy tools have in particular been applied to integrated membrane proteins (30, 31), such as interaction networks like oxidative phosphorylation (32–36), cell division processes (37–39) and protein translocation (40), along with bacterial cell motility (41–44). The tools can also probe the aqueous environment of cells as opposed to just on their hydrophobic cell membrane surface, including processes of DNA replication/remodeling/repair (45–47), and systems more directly relevant to biomedicine such as bacterial infection (48–50).

In this Points of View article I discuss further the findings from my team from single-molecule fluorescence microscopy to track functional TFs with very high speed to match

typical rates of protein diffusion in live cells and thereby enable ‘blur-free’ observations. We were able to quantify the composition and dynamics of Mig1 under normal and perturbed conditions which affected its state of phosphorylation, and also performed experiments on a protein called Msn2 which functions antagonistically, i.e. instead as an enhancer/activator, for many of the same Mig1 target genes (51) through a completely different signaling pathway. The results showed unexpectedly that Mig1 binds to its target genes as an oligomeric cluster which has stoichiometries in the range ~6-9 molecules. We found evidence that Mig1 molecules in a cluster are glued together through interactions of intrinsically disordered peptide sequences innervated by molecular crowding depletion forces in the cell. Our findings may reveal a more general eukaryotic cell strategy for the control of gene expression which uses intrinsic disorder of many TFs to form clusters that then enable large reductions in the time taken to find a given target gene.

## Results

### Single-molecule optical proteomics indicates the presence of Mig1 oligomeric clusters

We used millisecond Slimfield single-molecule fluorescence imaging (46, 52, 53) on live *S. cerevisiae* cells (Fig. 1A) using a green fluorescent protein (GFP) reporter for Mig1 integrated into the genome, including mCherry reporter on the RNA polymerase subunit protein Nrd1 to indicate the position of the cell nucleus. Slimfield was optimized for single-molecule detection sensitivity by using an *in vitro* imaging assay (54). We also measured the maturation effect of the fluorescent proteins in these cells (55) and estimate it to be <15% immature fluorescent protein over the timescale of imaging experiments. Note, Slimfield limits the observation area to an equivalent diameter of <10  $\mu\text{m}$  in the lateral plane to achieve rapid imaging sample times of millisecond and, in some instances, sub-millisecond levels (56), but is less ideal to eukaryotic imaging of cells with larger nuclei. A host of other single-molecule techniques based on light-sheet imaging have larger fields of view, and also combine low background and low light toxicity. For the interested reader, these include: HILO (by Tokunaga M.N. et al. (57)), AFM cantilever lightsheet (by Gebhardt, J.C. et al. (11)), lattice light-sheet (by Chen B.C. et al. (58)), multi-focus (by Abrahamsson S. et al. (59)), remote focusing (by Yang et al. (60)), and diagonally scanned light sheet (by Dean et al. (61)).

Under depleted /elevated extracellular *glucose* (-/+), we measured cytoplasmic and nuclear Mig1 localization bias respectively, visible in individual cells by our generating rapid microfluidic exchange (a few seconds) of extracellular fluid (Fig. 1B), and resolved two components under both conditions consistent with a diffuse monomeric pool and distinct oligomeric foci of Mig1 (Fig. 1C). The foci were also visible as hotspots using the green-red photoswitchable fluorescent protein mEos2 (62) excited by super-resolution stochastic optical reconstruction microscopy (STORM) (Fig. 1C), with modeling using 3C structural data of the yeast chromosome (63) and sequence alignment analysis for the location of Mig1 target promoters supporting the hypothesis that the majority of Mig1 clusters were specifically binding to Mig1 target genes.

Nanoscale tracking determined the position of tracked Mig1 foci to a lateral precision of 40 nm (33, 64) coupled to stoichiometry analysis using stepwise photobleaching of GFP (54) and single cell copy number analysis (65). An additional output from the tracking was the effective diffusion coefficient  $D$  as a function of its location in either the cytoplasm, nucleus or translocating across the nuclear envelope, as well as the copy number of Mig1 molecules associated with each subcellular region and in each cell as a whole, indicating ~850-1,300 Mig1 molecules per cell dependent on extracellular glucose. It should be noted

that confinement may affect the apparent diffusion coefficient in the small volume of a yeast nucleus if the length the mean square displacement (MSD) of tracked particles is comparable to the diameter of the nucleus, however, if our case only the short length scale MSD regions are considered to determine  $D$ .

In control experiments, a modified strain (51) generated with a binding site for protein PP7 on mRNA produced by one of the Mig1 target genes called *GSY1* showed colocalization between PP7-GFP expressed off a plasmid and Mig1-mCherry expressed genomically under high glucose conditions. We also observed similar clustering and co-localization to PP7 for the antagonistic TF Msn2. These PP7 co-localization results suggest that clusters both of Mig1 and Msn2 are *functionally* active in regulating target gene expression of the test target gene *GSY1*.

### **Cytoplasmic Mig1 diffuses rapidly but nuclear Mig1 can be mobile and immobile**

Cytoplasmic Mig1 fluorescent foci at *glucose* (+/-), and nuclear foci at *glucose* (-), were consistent with just a single mobile population whose  $D$  of 1-2  $\mu\text{m}^2/\text{s}$  consistent with earlier observations. However, nuclear foci at *glucose* (+) indicated a mixture of mobile and immobile components (Fig. 1D). These results suggested 20-30% of nuclear foci are immobile, consistent with a DNA-bound state. MSD analysis of foci tracks indicated Brownian diffusion over a few tens of ms but increasingly anomalous diffusion over longer timescales, consistent with *glucose* (+) Mig1 diffusion being impacted by interactions with nuclear structures, similar to that reported for other TFs (66). Here, this interaction depended on extracellular glucose despite Mig1 requiring a pathway of proteins to detect it, unlike the more direct detection mechanism of the prokaryotic *lac* repressor. Control experiments with Zn finger deletion strains of Mig1 indicated that Mig1 clusters bind to the DNA via their Zn finger motif with direct glucose dependence. At the high laser excitation intensities used for Slimfield imaging photobleaching is rapid, and so typically a single GFP molecule will photobleach on average after 5-10 consecutive image frame. To account for this we interpolate observed foci brightness values back to the start of each photobleach using an exponential photobleach function. We observed no direct evidence for irreversible photobleaching (i.e. 'photoblinking') with GFP at these intensities, though other fluorescent proteins such as YFP have been known to exhibit such blinking behavior, which if so would need to be further characterized, for example using surface immobilized purified YFP *in vitro* samples. A general compromise here, however, is to confine tracking analysis to typically less than 100 ms of laser exposure so that irreversible photoblinking is more dominant than reversible blinking.

### **Mig1 nuclear pore complex selectivity is mediated by interactions distant from the nuclear envelope**

We compared the spatiotemporal dynamics of different Mig1 clusters during translocation by converting trans-nuclear tracks into coordinates parallel and perpendicular to the measured nuclear envelope location, and synchronizing coordinate origins to be at the nuclear envelope crossing point for a given foci track. A heat map of spatial locations of translocating clusters indicated a hotspot of comparable volume to the nuclear pore complexes and accessory structures (67, 68) (Fig. 1E). The dwell time during nuclear envelope translocation was ~10 ms, similar to previous estimates for transport factors (69) but here found to be insensitive to glucose (Fig. 1F), demonstrating that there is no direct selectivity on the basis

of TF phosphorylation state by nuclear pore complexes themselves which suggests that cargo selectivity mechanisms of nuclear transport (70) are blind to phosphorylation state. Coupled with the observation that Mig1 at *glucose* (-) does not exhibit immobility in the nucleus and that Mig1 lacking the Zn finger still accumulates in the nucleus at *glucose* (+) this suggests that Mig1 localization is driven by changes in Mig1 binding affinity to other proteins, e.g. the general co-repressor complex at the genome (71), or outside the nucleus not involving the nuclear pore complex.

### **Mig1 nuclear clusters turn over in >100 s**

By modifying the microscope we were able to implement fluorescence recovery after photobleaching (FRAP) to probe nuclear turnover of Mig1, by focusing a separate laser onto just the nucleus, photobleaching this region with a rapid 200 ms pulse, and quantifying any subsequent fluorescence intensity recovery into that region (Fig. 1G). We could then acquire images with millisecond precision for individual frames but stroboscopically illuminating to extend the range of time scales for recovery before significant GFP photobleaching occurred, enabling FRAP observations at a single-molecule precision to timescales >1,000 s. Analyses demonstrated measurable recovery for both foci and the diffuse pool components in the nucleus, which could be fitted by single exponential functions indicating fast recovery of pool at both *glucose* (-) and (+) with a time constant of just a few seconds but a larger time constant at *glucose* (+) for nuclear foci of at least ~100s (Fig. 1H), with recovery of intensity being consistent with units of ~7-9 GFP molecules for the foci component but no obvious periodicity in stoichiometry measurable from pool recovery. These data suggested that molecular turnover at nuclear foci of Mig1 bound to target genes occurred in units of whole Mig1 clusters.

### **Clusters are stabilized by molecular crowding and intrinsic disorder**

Native, denaturing gel electrophoresis and western blots on purified extracts from Mig1-GFP cells (Fig. 1I) indicated a single band corresponding to Mig1. *In vitro* Slimfield imaging of purified Mig1-GFP under identical imaging conditions for live cells similarly indicated monomeric Mig1-GFP foci in addition to a small fraction of brighter foci which were consistent with chance overlap of monomer GFP images. However, addition of a molecular crowding reagent in the form of low molecular weight polyethylene glycol (PEG) at a concentration known to correspond to small molecule 'depletion' forces in cells (72) resulted in significant numbers of oligomers (Fig. 1J), suggesting that Mig1 clusters present in live cells regardless of glucose may be stabilized by depletion components that are lost during biochemical purification.

Secondary structure predictions suggested significant regions of disorder away from the Zn finger binding motif. We measured changes in circular dichroism of the Mig1 fusion construct upon addition of PEG (Fig. 1K) in a wavelength range known to be sensitive to transitions between ordered and intrinsically disordered states (73, 74). We also noted similar levels of disorder content in the Msn2 protein far from the Zn finger motif. These observations suggested a TF 'molecular bipolarity', in regards to disorder content, which stabilizes a cluster compact core focused around the disordered regions that undergo a putative phase transition to a more structure state, while exposing Zn fingers and positive surface charges to enable specific and non-specific interactions with accessible DNA strands (Fig. 1L).

## 210 **Perspective**

211 Our findings address aspects of functional gene regulation in live cells which have hitherto  
 212 been unexplored, using biophysical technology that has not been available until recently. The  
 213 results strongly support a functional link between Mig1 and Msn2 TF clusters and target gene  
 214 expression; a biological role of multivalent TFs for enhancing intersegmental transfer had  
 215 been elucidated previously in simulations (9) but unobserved experimentally until our  
 216 discoveries here, and so our findings impact on the longstanding question of how TFs might  
 217 find their targets in the genome so efficiently. Clustering of a range of nuclear factors has  
 218 been observed previously using single-molecule techniques, such as transient RNA  
 219 Polymerase II cluster dynamics in living cells using time-correlated PALM (tc-PALM) (75,  
 220 76). Also functional nuclear protein clusters have been seen (77) and the Bicoid transcription  
 221 factor in fruit fly embryos has been observed to form clusters mediated in part mediated by  
 222 intrinsically disordered peptide sequences (78).

223 Quantifying nearest-neighbor distances between Mig1 promoter sites in the *S.*  
 224 *cerevisiae* genome from the 3C model indicates 20-30% are <50 nm apart, small enough to  
 225 enable different DNA segments to be linked though intersegment transfer by a single cluster  
 226 (6, 9), which would also enable in principle simultaneous binding of >1 gene target from just  
 227 a single TF cluster. There is a net positive charge in the vicinity of Zn finger motifs, and this  
 228 would also enable non-specific electrostatic interactions with the negatively charged  
 229 phosphate backbone of DNA, facilitating 1D sliding diffusion of the protein along a DNA  
 230 strand. Thus, a cluster may be able to slide along DNA in a largely sequence-independent  
 231 manner and undergo intersegmental transfer to a neighboring strand relatively easily, either  
 232 spontaneously or stimulated by the presence of protein barriers on the DNA in a process  
 233 likely to have some sequence dependence when an obstacle is encountered. In particular,  
 234 bound RNA polymerases present during gene transcription at sequence specific sites could  
 235 act as roadblocks to kick off translocating clusters from a DNA strand, to again facilitate  
 236 intersegmental transfer and thus increase the ultimate chances that TF clusters will encounter  
 237 one of the gene targets and specifically bind via the Zn finger motif, thus predominantly  
 238 circumventing the requirement for significant amounts of slow 3D diffusion in the  
 239 nucleoplasm.

240 Our discovery is, to our knowledge, the first to make a link between predicted disorder  
 241 and the ability to form oligomeric clusters in TFs. Our findings may potentially offer some  
 242 insights into addressing the longstanding question of why in general there is so much  
 243 predicted disorder in eukaryote transcription factors; ~90% of eukaryotic TFs indicate  
 244 significant proportions of sequences with disordered content (79). Our finding that protein  
 245 interactions based on relatively weak molecular crowding depletion forces has functional  
 246 relevance in several areas of cell biology, such as processes involving aggregation mediated  
 247 through intrinsic disorder interactions; for example, those of amyloid plaques found in  
 248 neurodegenerative disorders including Alzheimer's and Parkinson's diseases (80). Increased  
 249 understanding of the clustering mechanism might therefore be of value in understanding the  
 250 progression of these diseases. Open questions remain though: for example, are clusters homo-  
 251 oligomeric or do they contain multiple different TFs? How is specificity maintained inside a  
 252 cluster? Are the components of the clusters themselves dynamic and undergo molecular  
 253 turnover? Can the ability to cluster be controlled, for example by switching the state of  
 254 phosphorylation?  
 255

## 256 **Acknowledgments**



The work of the original research article described (24) also involved Adam Wollman, Sviatlana Shashkova, Erik Hedlund, Rosemarie Friemann and Stefan Hohmann, and the Bioscience Technology Facility of the University of York, UK. Thanks to Mark Johnston (CU Denver) for a Mig1 phosphorylation mutant plasmid, and Michael Elowitz (Caltech) for a Mig1/Msn2/PP7 and Zn finger deletion strain.

## Funding

Supported by the Biological Physical Sciences Institute, Royal Society, MRC (grant MR/K01580X/1), BBSRC (grant BB/N006453/1), Swedish Research Council and European Commission via Marie Curie-Network for Initial training ISOLATE (Grant agreement nr: 289995) and the Marie Curie Alumni Association.

## Conflict of interest

There are no conflicts of interests.

## References

1. F. Jacob, J. Monod, Genetic regulatory mechanisms in the synthesis of proteins. *J. Mol. Biol.* **3**, 318–356 (1961).
2. J. Gertz, E. D. Siggia, B. A. Cohen, Analysis of combinatorial cis-regulation in synthetic and genomic promoters. *Nature*. **457**, 215–8 (2009).
3. O. G. Berg, R. B. Winter, P. H. Von Hippel, Diffusion-driven mechanisms of protein translocation on nucleic acids. 1. Models and theory. *Biochemistry*. **20**, 6929–6948 (1981).
4. A. Mahmutovic, O. G. Berg, J. Elf, What matters for lac repressor search in vivo--sliding, hopping, intersegment transfer, crowding on DNA or recognition? *Nucleic Acids Res.* **43**, 3454–64 (2015).
5. S. E. Halford, J. F. Marko, How do site-specific DNA-binding proteins find their targets? *Nucleic Acids Res.* **32**, 3040–52 (2004).
6. D. M. Gowers, S. E. Halford, Protein motion from non-specific to specific DNA by three-dimensional routes aided by supercoiling. *EMBO J.* **22**, 1410–8 (2003).
7. S. T. Whiteside, S. Goodbourn, Signal transduction and nuclear targeting: regulation of transcription factor activity by subcellular localisation. *J. Cell Sci.* **104** ( Pt 4, 949–55 (1993).
8. C. T. Harbison *et al.*, Transcriptional regulatory code of a eukaryotic genome. *Nature*. **431**, 99–104 (2004).
9. H. G. Schmidt, S. Sewitz, S. S. Andrews, K. Lipkow, An Integrated Model of Transcription Factor Diffusion Shows the Importance of Intersegmental Transfer and Quaternary Protein Structure for Target Site Finding. **9**, e108575 (2014).
10. G.-W. Li, X. S. Xie, Central dogma at the single-molecule level in living cells. *Nature*. **475**, 308–15 (2011).
11. J. C. M. Gebhardt *et al.*, Single-molecule imaging of transcription factor binding to DNA in live mammalian cells. *Nat. Methods*. **10**, 421–426 (2013).


12. D. Normanno *et al.*, Probing the target search of DNA-binding proteins in mammalian cells using TetR as model searcher. *Nat. Commun.* **6**, 7357 (2015).
13. D. Mazza, A. Abernathy, N. Golob, T. Morisaki, J. G. McNally, A benchmark for chromatin binding measurements in live cells. *Nucleic Acids Res.* **40**, e119 (2012).
14. Z. Liu *et al.*, 3D imaging of Sox2 enhancer clusters in embryonic stem cells. *Elife.* **3**, e04236 (2014).
15. J. Chen *et al.*, Single-Molecule Dynamics of Enhanceosome Assembly in Embryonic Stem Cells. *Cell.* **156**, 1274–1285 (2014).
16. Z. Zhang *et al.*, Rapid dynamics of general transcription factor TFIIB binding during preinitiation complex assembly revealed by single-molecule analysis. *Genes Dev.* **30**, 2106–2118 (2016).
17. P. Hammar *et al.*, The lac repressor displays facilitated diffusion in living cells. *Science.* **336**, 1595–8 (2012).
18. J. C. M. Gebhardt *et al.*, Single-molecule imaging of transcription factor binding to DNA in live mammalian cells. *Nat. Methods.* **10**, 421–6 (2013).
19. J. O. Nehlin, M. Carlberg, H. Ronne, Control of yeast GAL genes by MIG1 repressor: a transcriptional cascade in the glucose response. *EMBO J.* **10**, 3373–7 (1991).
20. E. Frolova, Binding of the glucose-dependent Mig1p repressor to the GAL1 and GAL4 promoters in vivo: regulation by glucose and chromatin structure. *Nucleic Acids Res.* **27**, 1350–1358 (1999).
21. M. J. De Vit, J. a Waddle, M. Johnston, Regulated nuclear translocation of the Mig1 glucose repressor. *Mol. Biol. Cell.* **8**, 1603–18 (1997).
22. L. Bendrioua *et al.*, Yeast AMP-activated Protein Kinase Monitors Glucose Concentration Changes and Absolute Glucose Levels. *J. Biol. Chem.* **289**, 12863–75 (2014).
23. S. Shashkova, A. J. M. Wollman, M. C. Leake, S. Hohmann, The yeast Mig1 transcriptional repressor is dephosphorylated by glucose-dependent and -independent mechanisms. *FEMS Microbiol. Lett.* **364** (2017), doi:10.1093/femsle/fnx133.
24. A. J. M. J. Wollman *et al.*, Transcription factor clusters regulate genes in eukaryotic cells. **6**, e27451 (2017).
25. M. C. Leake, The physics of life: one molecule at a time. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **368**, 20120248 (2013).
26. H. Miller, Z. Zhou, J. Shepherd, A. J. M. Wollman, M. C. Leake, Single-molecule techniques in biophysics: a review of the progress in methods and applications. *Reports Prog. Phys.* **81**, 24601 (2018).
27. S. Shashkova, M. C. Leake, Single-molecule fluorescence microscopy review: Shedding new light on old problems. *Biosci. Rep.* **37** (2017), doi:10.1042/BSR20170031.
28. B. Huang *et al.*, Counting Low-Copy Number Proteins in a Single Cell. *Science.* **315**, 81–4 (2007).
29. M. Wu, A. K. Singh, Single-cell protein analysis. *Curr. Opin. Biotechnol.* **23**, 83–8 (2012).

30. S. J. Bryan *et al.*, Localisation and interactions of the Vipp1 protein in cyanobacteria. *Mol. Microbiol.* **94**, 1179–1195 (2014).
31. A. Nenninger *et al.*, Independent mobility of proteins and lipids in the plasma membrane of *Escherichia coli*. *Mol. Microbiol.* **92**, 1142–53 (2014).
32. T. Lenn, M. C. Leake, Single-molecule studies of the dynamics and interactions of bacterial OXPHOS complexes. *Biochim. Biophys. Acta - Bioenerg.* **1857**, 224–231 (2016).
33. I. Llorente-Garcia *et al.*, Single-molecule in vivo imaging of bacterial respiratory complexes indicates delocalized oxidative phosphorylation. *Biochim. Biophys. Acta.* **1837**, 811–24 (2014).
34. Y. T. Lenn, M. C. Leake, C. W. Mullineaux, Are *Escherichia coli* OXPHOS complexes concentrated in specialized zones within the plasma membrane? *Biochem. Soc. Trans.* **36**, 1032–6 (2008).
35. T. Lenn, M. C. Leake, C. W. Mullineaux, Clustering and dynamics of cytochrome bd-I complexes in the *Escherichia coli* plasma membrane in vivo. *Mol. Microbiol.* **70**, 1397–407 (2008).
36. A. Badrinarayanan, M. C. Leake, Using Fluorescence Recovery After Photobleaching (FRAP) to Study Dynamics of the Structural Maintenance of Chromosome (SMC) Complex In Vivo. *Methods Mol. Biol.* **1431**, 37–46 (2016).
37. S.-W. S.-W. Chiu, M. A. J. Roberts, M. C. Leake, J. P. Armitage, Positioning of chemosensory proteins and ftsz through the *rhodobacter sphaeroides* cell cycle. *Mol. Microbiol.* **90**, 322–37 (2013).
38. A. W. Bisson-Filho *et al.*, Treadmilling by FtsZ filaments drives peptidoglycan synthesis and bacterial cell division. *Science.* **355**, 739–743 (2017).
39. V. A. Lund *et al.*, Molecular coordination of *Staphylococcus aureus* cell division. *Elife.* **7**, e32057 (2018).
40. M. C. Leake *et al.*, Variable stoichiometry of the TatA component of the twin-arginine protein transport system observed by in vivo single-molecule imaging. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 15376–81 (2008).
41. S. W. W. Reid *et al.*, The maximum number of torque-generating units in the flagellar motor of *Escherichia coli* is at least 11. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 8066–71 (2006).
42. Y. Sowa *et al.*, Direct observation of steps in rotation of the bacterial flagellar motor. *Nature.* **437**, 916–9 (2005).
43. T. Pilizota *et al.*, A molecular brake, not a clutch, stops the *Rhodobacter sphaeroides* flagellar motor. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 11582–7 (2009).
44. C.-J. Lo, M. C. Leake, T. Pilizota, R. M. Berry, Nonequivalence of membrane voltage and ion-gradient as driving forces for the bacterial flagellar motor at low load. *Biophys. J.* **93**, 294–302 (2007).
45. R. Reyes-Lamothe, D. J. Sherratt, M. C. Leake, Stoichiometry and architecture of active DNA replication machinery in *Escherichia coli*. *Science.* **328**, 498–501 (2010).
46. A. Badrinarayanan, R. Reyes-Lamothe, S. Uphoff, M. C. Leake, D. J. Sherratt, In vivo architecture and action of bacterial structural maintenance of chromosome proteins.

384 *Science*. **338**, 528–31 (2012).


- 385 47. A. J. M. Wollman, A. H. Syeda, P. McGlynn, M. C. Leake, Single-molecule  
386 observation of DNA replication repair pathways in *E. coli*. *Adv. Exp. Med. Biol.* **915**,  
387 5–16 (2016).
- 388 48. M. C. Leake, The Biophysics of Infection. *Adv. Exp. Med. Biol.* **915**, 1–3 (2016).
- 389 49. H. Miller, A. J. M. Wollman, M. C. Leake, Designing a Single-Molecule Biophysics  
390 Tool for Characterising DNA Damage for Techniques that Kill Infectious Pathogens  
391 Through DNA Damage Effects. *Adv. Exp. Med. Biol.* **915**, 115–27 (2016).
- 392 50. A. J. M. Wollman, H. Miller, S. Foster, M. C. Leake, An automated image analysis  
393 framework for segmentation and division plane detection of single live *Staphylococcus*  
394 *aureus* cells which can operate at millisecond sampling time scales using bespoke  
395 Slimfield microscopy. *Phys. Biol.* **5**, 55002 (2016).
- 396 51. Y. Lin, C. H. Sohn, C. K. Dalal, L. Cai, M. B. Elowitz, Combinatorial gene regulation  
397 by modulation of relative pulse timing. *Nature*. **527**, 54–58 (2015).
- 398 52. M. Plank, G. H. Wadhams, M. C. Leake, Millisecond timescale slimfield imaging and  
399 automated quantification of single fluorescent protein molecules for use in probing  
400 complex biological processes. *Integr. Biol. (Camb)*. **1**, 602–12 (2009).
- 401 53. R. Reyes-Lamothe, D. J. Sherratt, M. C. Leake, Stoichiometry and architecture of  
402 active DNA replication machinery in *Escherichia coli*. *Science*. **328**, 498–501 (2010).
- 403 54. M. C. Leake *et al.*, Stoichiometry and turnover in single, functioning membrane  
404 protein complexes. *Nature*. **443**, 355–358 (2006).
- 405 55. S. Shashkova, A. Wollman, S. Hohmann, M. C. Leake, Characterising Maturation of  
406 GFP and mCherry of Genomically Integrated Fusions in *Saccharomyces cerevisiae*.  
407 *Bio Protoc.* **7**, e2710 (2018).
- 408 56. H. Miller, J. Cosgrove, A. Wollman, E. Taylor, P. O’Toole, M. Coles, M. C. Leake.  
409 High-speed single-molecule tracking of CXCL13 in the B-Follicle. *Front. Immunol.*,  
410 in press (2018).
- 411 57. M. Tokunaga, N. Imamoto, K. Sakata-Sogawa, Highly inclined thin illumination  
412 enables clear single-molecule imaging in cells. *Nat. Methods*. **5**, 159–161 (2008).
- 413 58. B.-C. Chen *et al.*, Lattice light-sheet microscopy: imaging molecules to embryos at  
414 high spatiotemporal resolution. *Science*. **346**, 1257998 (2014).
- 415 59. S. Abrahamsson *et al.*, Fast multicolor 3D imaging using aberration-corrected  
416 multifocus microscopy. *Nat. Methods*. **10**, 60–63 (2013).
- 417 60. B. Yang *et al.*, High Numerical Aperture Epi-illumination Selective Plane Illumination  
418 Microscopy. *bioRxiv*, 273359 (2018).
- 419 61. K. M. Dean *et al.*, Diagonally Scanned Light-Sheet Microscopy for Fast Volumetric  
420 Imaging of Adherent Cells. *Biophys. J.* **110**, 1456–1465 (2016).
- 421 62. S. A. McKinney, C. S. Murphy, K. L. Hazelwood, M. W. Davidson, L. L. Looger, A  
422 bright and photostable photoconvertible fluorescent protein. *Nat. Methods*. **6**, 131–3  
423 (2009).
- 424 63. Z. Duan *et al.*, A three-dimensional model of the yeast genome. *Nature*. **465**, 363–7  
425 (2010).

64. H. Miller, Z. Zhou, A. J. M. Wollman, M. C. Leake, Superresolution imaging of single DNA molecules using stochastic photoblinking of minor groove and intercalating dyes. *Methods*. **88**, 81–8 (2015).
65. A. J. M. Wollman, M. C. Leake, Millisecond single-molecule localization microscopy combined with convolution analysis and automated image segmentation to determine protein concentrations in complexly structured, functional cells, one cell at a time. *Faraday Discuss.* **184**, 401–24 (2015).
66. I. Izeddin *et al.*, Single-molecule tracking in live cells reveals distinct target-search strategies of transcription factors in the nucleus. *Elife*. **3**, e02230 (2014).
67. S. A. Adam, The nuclear pore complex. *Genome Biol.* **2**, reviews0007.1-reviews0007.7 (2001).
68. C. Strambio-De-Castillia, M. Niepel, M. P. Rout, The nuclear pore complex: bridging nuclear transport and gene regulation. *Nat. Rev. Mol. Cell Biol.* **11**, 490–501 (2010).
69. W. Yang, J. Gelles, S. M. Musser, Imaging of single-molecule translocation through nuclear pore complexes. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 12887–12892 (2004).
70. A. R. Lowe *et al.*, Selectivity mechanism of the nuclear pore complex characterized by single cargo tracking. *Nature*. **467**, 600–603 (2010).
71. M. A. Treitel, M. Carlson, Repression by SSN6-TUP1 is directed by MIG1, a repressor/activator protein. *Proc. Natl. Acad. Sci. U. S. A.* **92**, 3132–6 (1995).
72. Y. Phillip, G. Schreiber, Formation of protein complexes in crowded environments--from in vitro to in vivo. *FEBS Lett.* **587**, 1046–52 (2013).
73. K. Sode, S. Ochiai, N. Kobayashi, E. Usuzaka, Effect of reparation of repeat sequences in the human alpha-synuclein on fibrillation ability. *Int. J. Biol. Sci.* **3**, 1–7 (2007).
74. C. Avitabile *et al.*, Circular Dichroism studies on the interactions of antimicrobial peptides with bacterial cells. *Sci. Rep.* **4**, 337–360 (2014).
75. I. I. Cisse *et al.*, Real-Time Dynamics of RNA Polymerase II Clustering in Live Human Cells. *Science*. **341**, 664–667 (2013).
76. W.-K. Cho *et al.*, RNA Polymerase II cluster dynamics predict mRNA output in living cells. *Elife*. **5**, e13617 (2016).
77. J. Qian *et al.*, B Cell Super-Enhancers and Regulatory Clusters Recruit AID Tumorigenic Activity. *Cell*. **159**, 1524–1537 (2014).
78. M. Mir *et al.*, Dense Bicoid Hubs Accentuate Binding along the Morphogen Gradient. *Genes & Dev.* **31**, 1784–94 (2017).
79. J. Liu *et al.*, Intrinsic disorder in transcription factors. *Biochemistry*. **45**, 6873–88 (2006).
80. V. N. Uversky, V. B. Patel, Intrinsically disordered proteins and their (disordered) proteomes in neurodegenerative disorders. *Front Aging Neurosci.* **7**, 18 (2015).
81. M. C. Leake, D. Wilson, M. Gautel, R. M. Simmons, The elasticity of single titin molecules using a two-bead optical tweezers assay. *Biophys. J.* **87**, 1112–35 (2004).
82. M. C. Leake, D. Wilson, B. Bullard, R. M. Simmons, The elasticity of single kettin molecules using a two-bead laser-tweezers assay. *FEBS Lett.* **535**, 55–60 (2003).



**Figure 1.** TFs form clusters in eukaryotic cell. (A) Schematic of millisecond Slimfield microscopy. (B) Fluorescence imaging of Mig1-GFP (green) with nucleus indicated (red) by Nrd1-mCherry, showing different cellular locations, stoichiometry determined by step-wise photobleaching that can be measured using Fourier analysis and edge-detection filters (54, 81, 82). (C) STORM imaging using Mig1-mEos2. (D) Mobility analysis for cumulative distribution function (CDF) and Gamma fits. (E) Mig-GFP localization through a nuclear pore complex. (F) Dwell time for tracks translocating the nuclear envelope. (G) Images and (H) analysis for FRAP indicating turnover of nuclear Mig1-GFP. (I) Native and denaturing gels on purified Mig1-GFP. (J) Mig1-GFP cluster stoichiometry in presence/absence of molecular crowding. (K) Circular dichroism spectra in presence/absence of molecular crowding. (L) Cartoon model for shape of a Mig1 cluster in vicinity of DNA strands.







Biological Physical Sciences Institute  
Physical and Life Sciences at the Cutting-Edge Interface  
Building bridges. Removing barriers.

From: Professor Mark C. Leake FInstP, FRMS, FRSB, PhD  
Tel: +44 (0) 1904 322697/328566  
Fax: +44 (0) 1904 322214  
Email: [mark.leake@york.ac.uk](mailto:mark.leake@york.ac.uk)  
Institute: <http://www.york.ac.uk/physics/bpsi>  
Group: <http://single-molecule-biophysics.org>

Director, Biological Physical Sciences Institute (BPSI)  
Chair of Biological Physics  
Departments of Physics and Biology  
University of York  
Heslington  
York YO10 5DD, UK

Monday, 30 April 2018

**In reference to:** Reviewer comments for “Transcription factors in eukaryotic cells can functionally regulate gene expression by acting in oligomeric assemblies formed from an intrinsically disordered protein phase transition enabled by molecular crowding”

**Alberto Kornblihtt**

Dear Alberto

Many thanks for supplying the reviewer feedback: they were very helpful. I have revised the piece accordingly, and it is improved as a result. Please find overleaf a point by point response to all the reviewer comments.

If I may be of any further assistance please do not hesitate to contact me.

Yours sincerely,

Prof. Mark C. Leake

.\*\*\*



Reviewer 1:

Here is a small list of omitted citations on this subject that should be included to make the review more fairly balanced:

GR (Gebhardt, J.C. et al. <https://www.nature.com/articles/nmeth.2411>)  
TetR (Normanno, D. et al. <https://www.nature.com/articles/ncomms8357>)  
p53 (Mazza, D. et al. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3424588/>)  
Sox2 (Liu Z. et al. <https://elifesciences.org/articles/04236>  
and  
Chen J. et al. <https://www.ncbi.nlm.nih.gov/pubmed/24630727>)  
TFIIB  
(by Zhang Z. et al. <http://genesdev.cshlp.org/content/30/18/2106>)

These references have now been added

2. The only imaging modality presented in this mini-review is Slimfield. This technique limits the observation area to a few micrometers, and hence is unsuitable to eukaryotic imaging of cells with larger nuclei. A host of other single-molecule techniques based on light-sheet imaging have much bigger fields of view, and have the added advantage of low background and low light toxicity. Here are omitted references of imaging modalities that have been recently implemented to image eukaryotic nuclei:

HILO  
(by Tokunaga M.N. et al. <https://www.nature.com/articles/nmeth1171>)  
AFM cantilever light-sheet  
(by Gebhardt, J.C. et al. <https://www.nature.com/articles/nmeth.2411>)  
lattice light-sheet  
(by Chen B.C. et al. <http://science.sciencemag.org/content/346/6208/1257998>)  
multi-focus  
(by Abrahamsson S. et al. <https://www.nature.com/articles/nmeth.2277>)  
remote focusing  
(by Yang et al. <https://www.biorxiv.org/content/early/2018/02/28/273359>)  
diagonally scanned light sheet  
(by Dean et al. <https://doi.org/10.1016/j.bpj.2016.01.029>)

These references have now been added

3. Paragraph lines 50-64. Confusing mix of references for different model organisms: yeast, multi-cellular, and bacterial)

I have now clarified that these references refer to a range of different model organisms

4. Paragraph lines 114-126. The review should highlight how confinement affects apparent diffusion coefficient in the small volume of a yeast nucleus.

This has now been added

5. Paragraph lines 128-144. The review should state how bleaching and dye-photophysics (blinking, dark-state transitions) are accounted for. This is especially relevant for Slimfield imaging conditions of high laser power densities.

Discussion has been added here concerning how blinking and bleaching dye photophysics are accounted for.

6. Line 194: References 63 and 64 do not in fact argue for the static "transcription factories". These papers rather should be cited for time-correlated PALM (tc-PALM), which accurately accounts for dye-photophysics, and describe live cell RNA Polymerase II cluster dynamics that are quite transient.

This has now been added

7. Lines 33-34: Eukaryotic TFs do not always fluctuate between cytoplasm and the nucleus.

This has now been corrected

8. Line 88: The author should address how <15 % fluorophore maturation is compatible with single-molecule counting.

The '<15%' refers to the immature fluorescent protein level – this has now been clarified